Formulas

discrete time growth:

- $N_T = N_0 \lambda^T$
- $\lambda = f + p$
- $\mathcal{R} = f/(1-p)$

continuous time growth:

- $N(t) = N(0) \exp(rt)$
- \bullet r = b d
- $\mathcal{R} = b/d$


structured growth:

- $\bullet \ \ell_x = p_1 \times p_2 \times \dots p_{x-1}$
- $\mathcal{R} = \sum \ell_r f_r$
- $\sum \ell_x f_x \lambda^{-x} = 1$
- $SAD(x) \propto \ell_x \lambda^{-x}$
- 1. Quebec is a sprawling province, whose area exceeds that of the next-largest province (Ontario) by around 40 Mha. PEI is a small province with an area of < 1 Mha only about one tenth the area of the next smallest province. Which of these provinces is more unusual (more different from the others)?
 - A. Quebec is more unusual, whether we think additively or multiplicatively
 - $\mathbf{B.}$ PEI is more unusual, whether we think additively or multiplicatively
- C. Quebec is more unusual if we think additively, and PEI is more unusual if we thinking multiplicatively
- **D.** Quebec is more unusual if we think multiplicatively, and PEI is more unusual if we think additively
- **2.** What happens in the long run to a structured, unregulated population with *non-interacting* cohorts?
 - A. It reaches a stable age distribution
 - ${f B.}$ It reaches a stable population size
- C. The age distribution cycles while the total population size grows approximately exponentially
- **D.** The age distribution grows approximately exponentially while the total population size cycles

3. In a population of small mammals both sexes have a survival probability of 1/3, and females produce on average 3 offspring per year who survive to be counted. (both of these values are independent of population density). Which of the following is true?

- **A.** f = 3
- **B.** p = 2/3
- C. the average lifespan is 3 years
- **D.** the finite rate of growth λ depends on the sex ratio
- **E.** the finite rate of growth λ is 3.33
- **4.** Which of the following traits is *not* typically associated with r strategies?
 - **A.** Fast life cycle
 - **B.** Efficient resource use
 - C. Relatively small investment per offspring
 - **D.** Relatively large investment in dispersal
- 5. If we compare the stable age distribution from a relatively stable, developed country (eg., Sweden) and a rapidly growing, developing country (eg., Honduras), we expect to see that the growing country's age distribution is _____ than the stable country's.
 - **A.** more elliptical (with a wide middle)
 - **B.** less elliptical
 - C. more triangular (with a wide base)
 - **D.** less triangular
- **6.** Choose the most precise correct answer. For a population to be regulated, its percapita reproductive number must respond ______ to _____.
 - **A.** directly; population size
 - **B.** either directly or indirectly; population size
 - C. directly; population size or external factors like climate
 - **D.** either directly or indirectly; population size or external factors like climate
- 7. Choose the strongest correct answer. Dushoff believes that all ______ populations experience regulation _____
 - **A.** successful; all the time
 - **B.** successful; in the long term
 - C. surviving; all the time
 - **D.** surviving; in the long term


- **8.** Which of the following does *not* describe a life-history tradeoff?
- **A.** Organisms that use more resources for survival can reproduce more times, and produce more total offspring under good conditions
- **B.** Organisms that reproduce only once can produce more offspring at that time than if they saved resources so that they could survive and reproduce again
- C. Organisms that use more resources to help their offspring disperse cannot produce as many total offspring
- **D.** Organisms that use more resources to help their offspring survive cannot produce as many total offspring
- **9.** A group of tree species shows a tradeoff between r_{max} and K. We would expect to see:
- **A.** Species with high K doing well in less disturbed areas and species with high r_{max} doing well in more disturbed areas
- **B.** Species with high r_{max} doing well in less disturbed areas and species with high K doing well in more disturbed areas
 - C. Species with high K doing well in all areas
 - **D.** Species with high r_{max} doing well in all areas



Use the picture above for the following 2 questions.

10. What does this picture of survivorship in an idealized age-structured population indicate about *mortality* in this population?

- **A.** Mortality is constant
- **B.** Mortality is elevated in older individuals
- C. Mortality is elevated in younger individuals
- **D.** Mortality is elevated in both older and younger individuals
- **11.** The pictures below show *cumulative* survival. Which one corresponds to the picture shown above?



- 12. A certain species of moth reproduces only once in its lifetime. A researcher estimates that female moths that survive to become reproductively mature produce 150 eggs on average; that 40% of eggs will hatch; and that 3% of the female eggs that hatch will survive to become reproductively mature moths. If the moths lay equal numbers of male and female eggs, what is the finite rate of growth λ for this population? Assume that the time step of the model is equal to the amount of time it takes the moths to complete one generation (the generation time)
 - **A.** 0.45
 - **B.** 0.9
 - **C.** 1.8
 - **D.** 2.25

- **E.** 4.5
- 13. Given the information in the previous question, what is the strongest statement you can make about the reproductive number \mathcal{R} for this population?
 - **A.** λ and \mathcal{R} are both on the same side of 1
 - **B.** λ and \mathcal{R} are both on the same side of 0
 - C. \mathcal{R} is between 1 and λ
 - **D.** λ is between 1 and \mathcal{R}
 - **E.** λ and \mathcal{R} are equal
- 14. In some populations (like cod fish), it is hard to estimate the contribution of old individuals to population growth because old individuals have:
 - **A.** high values of both p and f
 - **B.** low values of p and high values of f
 - C. high values of both ℓ and f
 - **D.** low values of ℓ and high values of f
- 15. The size of a population of introduced beetles is estimated at 100 in 2014, and 200 in 2016. If we assume this population is growing exponentially, what population size would you predict for 2020?
 - **A.** 300
 - **B.** 400
 - C. 600
 - **D.** 800
 - **E.** 1600
- 16. Many species maximize their long-term average value of λ by:
 - A. speeding up their life cycle in general
 - **B.** slowing down their life cycle in general
 - C. speeding up their life cycle when conditions are good
 - **D.** slowing down their life cycle when conditions are good

Answer questions on this page in pen. Briefly show necessary work and equations. Points may be deducted for wrong information, even when the correct information is also there.

- 17. A species of plant produces 15 seeds on average in the first year after it is born, and 50 seeds on average in the second year after it is born, assuming it survives. Seeds survive the first year (and become adults) with probability 0.05, and first-year adults survive to become second-year adults with probability 0.7. Second-year adults always die.
- a) (2 points). If we model this population by counting *after* reproduction, how many age classes do we have? How would you describe the individuals censused in each age class, and how would you number the age classes?
- b) (1 points). How would this differ if we counted before reproduction?
- c) (2 points). If a scientist constructs life tables for the two scenarios above, what factors would you expect to be the same, and which ones could differ?
- d) (4 points). Construct a life table for the case where we count after reproduction (not the most common way). Show how you calculate the values of f_x , and then complete the life table below.

\boldsymbol{x}	$\int f_x$	p_x	ℓ_x	$\ell_x f_x$
1				
2				
$\overline{\mathcal{R}}$				